

Oral presentation

Open Access

## **Oxidative stress induces CHIP-mediated ubiquitination and proteasomal degradation of soluble guanylyl cyclase**

Sabine Meurer<sup>1,2</sup>, Tatjana Pabst<sup>1</sup>, Sylke Pioch<sup>1</sup>, Nils Opitz<sup>1,2</sup>, Peter M Schmidt<sup>2</sup>, Kristina Wagner<sup>1</sup>, Simone Matt<sup>1</sup>, Harald HHW Schmidt<sup>2</sup> and Werner Müller-Esterl<sup>\* 1</sup>

Address: <sup>1</sup>Institute of Biochemistry II, University of Frankfurt Medical School, Theodor-Stern-Kai 7, Frankfurt, Germany and <sup>2</sup>Department of Pharmacology & Centre for Vascular Health, Monash University, Clayton, Victoria, Australia

Email: Werner Müller-Esterl<sup>\*</sup> - [wme@biochem2.de](mailto:wme@biochem2.de)

<sup>\*</sup> Corresponding author

from 3<sup>rd</sup> International Conference on cGMP Generators, Effectors and Therapeutic Implications  
Dresden, Germany. 15–17 June 2007

Published: 25 July 2007

*BMC Pharmacology* 2007, **7**(Suppl 1):S34 doi:10.1186/1471-2210-7-S1-S34

This abstract is available from: <http://www.biomedcentral.com/1471-2210/7/S1/S34>

© 2007 Meurer et al; licensee BioMed Central Ltd.

Oxidative stress attenuates the NO-cGMP pathway, e.g. in the vascular system, through scavenging of free NO radicals by superoxide  $O_2^{\bullet-}$ , by inactivation of soluble guanylyl cyclase (sGC) via oxidation of its central  $Fe^{2+}$  ion, and by down-regulation of sGC protein levels. While the former pathways are well established, the molecular mechanisms underlying the latter are still obscure. Using oxidative sGC inhibitor ODQ we demonstrate rapid down-regulation of sGC protein in mammalian cells. Co-incubation with proteasomal inhibitor MG132 results in accumulation of ubiquitinated sGC whereas sGC activator BAY 58–2667 prevents ubiquitination. ODQ-induced down-regulation of sGC is mediated through selective ubiquitination of its b subunit, and BAY 58–2667 abrogates this effect. Ubiquitination of sGC-b is dramatically enhanced by E3 ligase CHIP. Our data indicate that oxidative stress promotes ubiquitination of sGC b subunit through E3 ligase CHIP, and that sGC activator 58–2667 reverts this effect, most likely through stabilization of the heme-free b subunit. Thus the deleterious effects of oxidative stress can be counter-balanced by an activator of a key enzyme of vascular homeostasis.